



Review Article

ISSN: 2454-5023
J. Ayu. Herb. Med.
2018; 4(1): 35-42
© 2018, All rights reserved
www.ayurvedjournal.com
Received: 03-02-2018
Accepted: 03-04-2018

Antidiabetic Potency of Bangladeshi Medicinal Plants

Towhid Hasan¹, Marjia Sultana²

¹ Department of Nutrition and Food Technology, Jessore University of Science and Technology, Jessore-7408, Bangladesh

² Department of Food Technology and Nutrition Science, Noakhali Science and Technology University, Noakhali-3814, Bangladesh

ABSTRACT

Diabetes mellitus is a degenerative disease being responsible for about 1.5 million deaths globally. In Bangladesh, the stress of diabetes is on rising and resulting in serious health implications along with significant economic crisis. Due to undesirable and inherent side effects, researchers are now shifting from the conventional therapy and trying to prevent and manage diabetes through traditional medicine. World Health Organization (WHO) also recommends the practice of customary herbal medicine for diabetes management, and support and encourage the augmentation of research to evaluate the hypoglycemic properties of the diverse medicinal plant species. Consequently, in the current review, the antihyperglycemic potency of some Bangladeshi medicinal plants has been evaluated and verified utilizing human as well as experimental animals. The results elucidate the glucose-lowering effects of the plants via different cellular mechanisms, including restoration of pancreatic β -cell, controlling the action of carbohydrate metabolizing enzymes, enhancing peripheral glucose utilization, increase in muscle glycogen store as well as activation of the insulin signaling cascade. In summary, this work may invigorate the researchers for more specific and focused research to provide a better and broad understanding of the antihyperglycemic mechanism and can act as an effective tool for choosing the plants with robust potential for unbolting of novel antidiabetic agents.

Keywords: Diabetes mellitus; antidiabetic medicinal plants; bioactive compounds; hypoglycemic potency; Bangladesh.

INTRODUCTION

Diabetes mellitus (DM) is a chronic, degenerative disease expressed by hyperglycemia (exalted blood glucose levels) arising from inherited and/or attained scarcity in insulin production from β islet cells of the pancreas, or by the incompetence of the insulin made [1]. Over 400 million people in the world are living with DM while an accounted 1.5 million deaths are attributed to it. In the past 35 years, the outbreak of DM has almost increased by twofold globally, rising from 4.7% to 8.5%. Epidemiologic studies have indicated that low- and middle-income countries have experienced a greater percentage of morbidity and mortality due to DM than high-income generating countries [2]. The World Health Organization (WHO) projected that, by 2030, DM is going to be one of the seven leading causes of death if the ongoing trend continues [3].

Diabetes may be classified based on the etiology and clinical symptoms as type 1 (also called insulin dependent diabetes mellitus), type 2 (also called non-insulin dependent diabetes mellitus), and gestational diabetes (a temporary condition during pregnancy). Being responsible for 90-95% of all cases, type 2 DM is the most prevalent one among diabetes. Diabetes of all types, if not controlled, has miserable upshot for health and well-being, and can lead to complications in many parts of the body causing serious damage to different organs, especially the eyes, kidneys, nerves, heart, and blood vessels, and can increase the overall risk of premature death [1]. Bangladesh, a lower-middle income country of South Asia, is stressed with the dual load of malnutrition. Non-communicable diseases like diabetes are on rising and resulting in serious health implications along with significant financial burden. Based on the recent findings, there has been a substantial increase in diabetes among Bangladeshi adults, from 4% in 2000 to 9.7% in 2014. More than half of the population with DM, 56%, are not aware that they have diabetes and don't receive any treatment [4].

Still now, diabetes management is a matter of concern worldwide, and effectual remedy has not been unfolded yet. There are copious therapeutic measurements attainable nowadays for diabetes comprising insulin and a number of oral antidiabetic medicines such as amylin analogs, α -glycosidase inhibitors include acarbose, miglitol and voglibose, sulfonylureas, biguanides, but it is a fact that full recuperation from diabetes, even of a single person, has never been proclaimed. The treatment of diabetes with synthetic drugs (oral hypoglycemic agents) are generally not promoted because of its high charge, and

*Corresponding author:

Marjia Sultana

Department of Food Technology
and Nutrition Science, Noakhali
Science and Technology
University, Noakhali-3814,
Bangladesh

Email: marjia28sultana[at]gmail.com

undesirable and inherent side effects such as hypoglycemia at higher doses, weight gain, liver problems, disturbances in the gastrointestinal tract (GIT), nausea, and diarrhea [5, 6]. Thus, the exploration of more efficient and safer therapy has continued to be a vital ground for investigation. Except for the prevailing therapeutic choices, abundant traditional herbal medicines have been advised for the amendment of diabetes and other non-communicable diseases. The traditional medicines have demonstrated their potentiality and showed an effulgent future in the therapy of diabetes and therefore, WHO recommends medicinal plants to be used for the rectification of DM [7, 8]. Herbal drugs are often prescribed owing to their efficacy, relatively low cost, and possession of little or no adverse effects [9].

Bangladesh, owing to its location and propitious climate, has been graced with an unbounded diversity of flora, together with medicinal plants. Among 5700 angiosperms growing in the country, above 500 are assumed to have medicinal properties [10, 11]. Being easily accessible and inexpensive, these natural herbal medicines have become an indispensable segment of public health services in Bangladesh. Hence, in this review, an effort was made to focus some experimentally ascertained antidiabetic medicinal plants of Bangladesh, and the bioactive compounds present and the probable mode of action through which they exert hypoglycemic potency.

METHODS

In the existing review, some ethnobotanical and ethnomedicinal surveys conducted in different districts and among different

communities of Bangladesh were utilized as a basis for the preparation of a list of medicinal plants commonly practiced to prevent and manage diabetes [12-15]. Afterward, electronic databases including Google Scholar, PubMed, and Cochrane library etc. were explored (till December 2017) for each and every medicinal plant, and all collected papers were reconnoitered to ensure any *in vivo*, *in vitro*, or clinical indication for their competence and pharmacological mechanisms of action. For the present study, published articles were considered solely eligible.

Medicinal plants with antidiabetic potency

In Bangladesh, traditional practices in diabetes treatment using medicinal plants in an unexplored horizon. Though, investigations of medicinal plants with various species were studied in this review (Table 1). Multiple segments of the plants (i.e. leaf, fruit, root, bark, bulb etc.) were used to carry out the studies. The initial phytochemical analysis illustrated the existence of flavonoids, terpenoids, saponins, alkaloids etc. Streptozotocin or alloxan-induced diabetic modules were employed to conduct the efficacy evaluation of the plants. The research results illuminated the glucose-lowering effects while proposing numerous mechanisms of action for the plant extracts which include the resurrection of pancreatic β -cell, regulation of the activity of carbohydrate metabolizing enzymes, increase in peripheral glucose utilization, increase in muscle glycogen store as well as activation of the insulin signaling cascade.

Table 1: Medicinal plants of Bangladesh with antidiabetic activity

Botanical name (Common Name)	Part(s) used	Bioactive constituent(s)	Probable mechanisms of action
<i>Acacia catechu</i> (L.f.) Willd. (Khair) [16]	▪ Bark	▪ Flavonoids	<ul style="list-style-type: none"> ▪ Stimulate insulin secretion ▪ Promote β-cell regeneration
<i>Adhatoda vasica</i> Nees (Basak) [17]	▪ Leaf	▪ Alkaloids	<ul style="list-style-type: none"> ▪ Inhibit activities of α-glucosidase
<i>Aegle marmelos</i> (L.) Corr. (Bel) [18, 19]	▪ Fruit ▪ Bark	▪ Tannins	<ul style="list-style-type: none"> ▪ Increase insulin secretion from pancreatic β-cell ▪ Regulate the activity of carbohydrate metabolizing enzymes
<i>Allium cepa</i> L. (Piaj) [20, 21]	▪ Bulb	<ul style="list-style-type: none"> ▪ Sulphur containing amino acids ▪ Flavonoids 	<ul style="list-style-type: none"> ▪ Control glucose metabolizing enzymes activities ▪ Upregulate expressions of insulin receptor and glucose transporter
<i>Allium sativum</i> L. (Rasun) [22, 23]	▪ Bulb	▪ Sulphur containing amino acids	<ul style="list-style-type: none"> ▪ Regulate the activities of glucose metabolizing enzymes ▪ Increase insulin secretion of from pancreatic β-cell
<i>Alocasia macrorrhizos</i> (L.) G. Don [24]	▪ Rhizome	▪ Flavonoids	<ul style="list-style-type: none"> ▪ Stimulate insulin secretion by pancreatic β-cells
<i>Aloe vera</i> (L.) Burm. f. (Ghritakumari) [25, 26]	▪ Leaf	▪ Phytosterols	<ul style="list-style-type: none"> ▪ Control the activity of the carbohydrate metabolizing enzymes ▪ Stimulate synthesis and/or release of insulin
<i>Amaranthus spinosus</i> L. (Katanotey) [27, 28]	▪ Leaf	<ul style="list-style-type: none"> ▪ Alkaloids ▪ Flavonoids 	<ul style="list-style-type: none"> ▪ Inhibit α-amylase activity
<i>Amaranthus viridis</i> L. (Notey) [29, 30]	▪ Leaf	<ul style="list-style-type: none"> ▪ Flavonoids ▪ Terpenoids ▪ Saponins ▪ Alkaloids 	<ul style="list-style-type: none"> ▪ Enhance insulin from pancreatic β-cells ▪ Inhibit the activity of α-amylase ▪ Increase glucose uptake by diaphragm ▪ Enhance glycogenesis by liver
<i>Amorphophallus paeoniifolius</i> (Dennst.) Nicolson [31]	▪ Corm	<ul style="list-style-type: none"> ▪ Polyphenols ▪ Flavonoids 	<ul style="list-style-type: none"> ▪ Alleviate intestinal and renal disaccharides
<i>Ananas comosus</i> (L.) Merr. (Anaras) [32, 33]	▪ Leaf	<ul style="list-style-type: none"> ▪ Alkaloids ▪ Flavonoids ▪ Tannins 	<ul style="list-style-type: none"> ▪ Enhance insulin sensitivity ▪ Regulate the activities of glycogen metabolizing enzymes
<i>Andrographis paniculata</i> (Burm.f.) Nees (Kalomegh) [34, 35]	▪ Leaf	▪ Flavonoids	<ul style="list-style-type: none"> ▪ Inhibit activities of α-amylase and α-glucosidase ▪ Reduce the activities of glucose-6-phosphatase
<i>Areca catechu</i> L. (Supari) [36]	▪ Areca nut	▪ Flavonoids	<ul style="list-style-type: none"> ▪ Inhibit activities of α-glucosidase
<i>Asparagus racemosus</i> Willd. (Satamuli) [37]	▪ Root	▪ Flavonoids	<ul style="list-style-type: none"> ▪ Inhibition carbohydrate digestion and absorption ▪ Enhance insulin secretion and action
<i>Azadirachta indica</i> A. Juss. (Neem) [38-40]	▪ Leaf	<ul style="list-style-type: none"> ▪ Flavonoids ▪ Alkaloids ▪ Tannins ▪ Saponins 	<ul style="list-style-type: none"> ▪ Improve the insulin signaling molecules ▪ Increase glucose utilization in the skeletal muscle ▪ Increase release of insulin from β-cells

<i>Barringtonia acutangula</i> (L.) Gaertn. (Hijal) [41, 42]	▪ Fruit	▪ Flavonoids ▪ Terpenoids ▪ Tannins ▪ Saponins ▪ Alkaloids	▪ Regenerate β -cells of the pancreas
<i>Basella rubra</i> L. (Puishak) [43]	▪ Leaf	▪ Flavonoids ▪ Terpenoids ▪ Tannins ▪ Saponins ▪ Alkaloids	▪ Inhibit activities of α -amylase and α -glucosidase
<i>Boerhaavia diffusa</i> L. (Punarnava) [44]	▪ Leaf	▪ Alkaloids	▪ Reduce the activities of gluconeogenic enzymes
<i>Bombax ceiba</i> L. (Shimul) [45]	▪ Thalamus	▪ Tannins	▪ Inhibit activities of α -amylase and α -glucosidase
<i>Butea Monosperma</i> (Lam.) Taub. (Palas) [46, 47]	▪ Flower	▪ Flavonoids ▪ Saponins ▪ Phytosterols	▪ Increase peripheral glucose utilization ▪ Stimulate insulin secretion
<i>Camellia sinensis</i> (L.) Kuntze (Cha) [48]	▪ Leaf ▪ Flower	▪ Alkaloids ▪ Flavonoids ▪ Tannins	▪ Inhibit activities of α -amylase and α -glucosidase
<i>Cardiospermum halicacabum</i> L. (Phutka) [49]	▪ Leaf	▪ Flavonoids ▪ Saponins	▪ Reduce the activity of glucokinase ▪ Elevate the activities of gluconeogenic enzymes ▪ Stimulate insulin secretion from pancreatic β -cells
<i>Carica papaya</i> L. (Pepe) [50]	▪ Leaf	▪ Alkaloids ▪ Flavonoid ▪ Saponins ▪ Tannins	
<i>Catharanthus roseus</i> (L.) G. Don (Nayantara) [51-53]	▪ Leaf ▪ Twig	▪ Alkaloids ▪ Flavonoids ▪ Tannins ▪ Saponins	▪ Increase utilization of glucose by liver ▪ Alter the activities of glucose metabolizing enzymes
<i>Centella asiatica</i> (L.) Urb. (Thankuni) [54, 55]	▪ Whole plant	▪ Phenolics ▪ Triterpenoids ▪ Flavonoids ▪ Essential oils	▪ Inhibit the activity of α -amylase ▪ Inhibit intestinal disaccharidase activity
<i>Cinnamomum tamala</i> (Buch.-Ham.) T. Nees & Eberm. (Tejpata) [56, 57]	▪ Bark ▪ Leaf		▪ Inhibit α -amylase activity ▪ Promote peripheral glucose utilization ▪ Increase muscle glycogen store ▪ Regulate the activity of carbohydrate metabolizing enzymes
<i>Cinnamomum zeylanicum</i> Blume (Daruchini) [58]	▪ Bark	▪ Essential oils	
<i>Clitoria ternatea</i> L. (Aparajita) [59, 60]	▪ Leaf	▪ Flavonoids	▪ Reduce the activities of glucose-6-phosphatase, and increase the activity of glucokinase ▪ Regenerate β -cells ▪ Regenerate β -cells ▪ Inhibit α -amylase activity
<i>Coccinia grandis</i> (L.) Voigt (Telakucha) [61, 62]	▪ Leaf	▪ Alkaloids ▪ Flavonoids ▪ Tannins	▪ Inhibit activities of α -amylase and α -glucosidase
<i>Curcuma longa</i> L. (Halud) [63]	▪ Rhizome	▪ Volatile oils	
<i>Datura metel</i> L. (Dhutura) [64]	▪ Seed	▪ Alkaloids	▪ Enhance secretion of insulin
<i>Dillenia indica</i> L. (Chalta) [65]	▪ Leaf	▪ Phytosterols ▪ Polyphenols ▪ Triterpenoids	▪ Inhibit α -amylase and α -glucosidase activities
<i>Ficus benghalensis</i> L. (Bot) [66]	▪ Bark	▪ Leucopelargonin	▪ Stimulate insulin secretion
<i>Foeniculum vulgare</i> Mill. (Mouri) [67]	▪ Seed	▪ Alkaloids ▪ Flavonoid ▪ Saponins ▪ Tannins	▪ Enhance insulin secretion ▪ Increase liver and kidney hexokinase level
<i>Hibiscus rosa-sinensis</i> L. (Jaba) [68]	▪ Flower	▪ Alkaloids ▪ Tannins ▪ Flavonoids	▪ Regenerate β -cells
<i>Ipomoea aquatica</i> Forssk. (Misti Alu) [69]	▪ Plant	▪ Phenolics ▪ Flavonoid ▪ Alkaloids ▪ Saponins	▪ Preserve islets area and cell population
<i>Jatropha curcas</i> L. (Jamalgota) [70]	▪ Leaf	▪ Flavonoids	▪ Regenerate and protect β -cells ▪ Inhibit α -amylase activity
<i>Kalanchoe pinnata</i> (Lam.) Pers. (Patharkuchi) [71-73]	▪ Leaf	▪ Flavonoids ▪ Triterpenoids	▪ Stimulate insulin secretion ▪ Increase the number of pancreatic β -cells ▪ Inhibit α -amylase activity
<i>Lagenaria siceraria</i> (Molina) Standl. (Lau/Kadu) [74]	▪ Pulp ▪ Seed	▪ Flavonoids ▪ Alkaloids ▪ Saponins	▪ Increase insulin production and secretion
<i>Lagerstroemia speciosa</i> (L.) Pers. (Jarul) [75]	▪ Leaf	▪ Tannins	▪ Suppress gluconeogenesis ▪ Stimulate glucose oxidation

<i>Lawsonia inermis</i> L. (Mehedi/Mendi) [76]	▪ Leaf	▪ Flavonoids ▪ Saponins ▪ Tannins ▪ Fixed oil	▪ Inhibit α -amylase activity
<i>Linum usitatissimum</i> L. (Tisi) [77]	▪ Seed		▪ Prevent damage in pancreatic β -cells
<i>Luffa acutangula</i> (L.) Roxb. (Jhinga) [78]	▪ Fruit	▪ Saponins ▪ Phenols ▪ Flavonoids	▪ Inhibit the activities α -glucosidase
<i>Luffa aegyptiaca</i> Mill. (Dhunda/Dhundul) [79]	▪ Leaf	▪ Steroids ▪ Alkaloids ▪ Saponins ▪ Tannins ▪ Saponins ▪ Sterols	▪ Increase secretion of insulin from pancreatic β -cells ▪ Inhibit α -amylase and α -glucosidase activities
<i>Madhuca indica</i> J. F. Gmel. (Mahua) [80]	▪ Bark	▪ Saponins ▪ Sterols ▪ Mangiferin	▪ Inhibit the activities of α -amylase and α -glucosidase
<i>Mangifera indica</i> L. (Aam) [81]	▪ Leaf		▪ Inhibit activities of α -amylase and α -glucosidase
<i>Nicotiana tabacum</i> L. (Tamak) [82]	▪ Leaf	▪ Flavonoids ▪ Tannins ▪ Terpenoid	
<i>Nigella sativa</i> L. (Kalojira/Kalijira) [83-85]	▪ Seed	▪ Polyphenols	▪ Increase insulin secretion ▪ Induce proliferation of pancreatic β -cells ▪ Stimulate glucose uptake in muscle and fat cells ▪ Inhibit activities of α -glucosidase ▪ Regulate the activity of glucose metabolizing enzymes ▪ Stimulate insulin secretion ▪ Regulate the activity of carbohydrate metabolizing enzymes ▪ Inhibit activities of α -amylase and α -glucosidase ▪ Stimulate glucose uptake ▪ Increase secretion of insulin
<i>Nymphaea stellata</i> Willd. (Sapla) [86, 87]	▪ Flower	▪ Phenolics	
<i>Ocimum sanctum</i> L. (Tulsi) [40, 88, 89]	▪ Leaf	▪ Flavonoids ▪ Phenolics	
<i>Oryza sativa</i> L. (Dhan) [90]	▪ Bran	▪ Phenolics ▪ Flavonoids	
<i>Pandanus fascicularis</i> Lam. (Keora) [91]	▪ Root	▪ Flavonoids ▪ Phenolics	
<i>Phyllanthus emblica</i> L. (Amla/Amlaki) [92]	▪ Fruit	▪ Flavonoids ▪ Tannins ▪ Terpenoid ▪ Alkaloids	▪ Inhibit the activities of disaccharidase ▪ Reduce intestinal glucose absorption
<i>Piper betle</i> L. (Pan) [93]	▪ Leaf	▪ Alkaloids ▪ Saponins ▪ Tannins ▪ Alkaloids	▪ Control glucose metabolizing enzyme activities
<i>Polyalthia longifolia</i> (Sonn.) Thwaites (Debdaru) [94]	▪ Leaf		▪ Inhibit α -amylase and α -glucosidase enzymes activity
<i>Punica granatum</i> L. (Dalim/Bedana) [95]	▪ Fruit	▪ Tannins ▪ Flavonoids ▪ Pectins	▪ Regenerate β -cells of pancreas
<i>Saraca asoca</i> (Roxb.) Willd. (Ashok) [96, 97]	▪ Leaf ▪ Flower	▪ Phytoosterols ▪ Flavonoids	▪ Increase pancreatic secretion of insulin from the β -cells
<i>Solanum melongena</i> L. (Begun) [98]	▪ Fruit	▪ Phenolics	▪ Inhibit activities of α -amylase and α -glucosidase
<i>Spinacia oleracea</i> L. (Palong shak) [99]	▪ Leaf	▪ Flavonoids	▪ Regenerate β -cells of pancreas
<i>Spondias pinnata</i> (L. f.) Kurz (Amra) [100]	▪ Bark	▪ Flavonoids	▪ Increase the biosynthesis of insulin in the pancreas ▪ Regenerate β -cells
<i>Syzygium cumini</i> (L.) Skeels (Jam) [101]	▪ Seed	▪ Tannins ▪ Flavonoids ▪ Phenolics	▪ Inhibit activities of α -glucosidase
<i>Tagetes erecta</i> L. (Genda) [102]	▪ Inflorescence	▪ Flavonoids	▪ Inhibit activities of α -glucosidase and α -amylase
<i>Tamarindus indica</i> L. (Tetul) [103-105]	▪ Seed	▪ Flavonoids	▪ Suppress glucose-6-phosphatase activity ▪ Restore pancreatic β -cells ▪ Inhibit intestinal glucose absorption ▪ Regenerate β -cells
<i>Tectona grandis</i> L. f. (Segun) [106]	▪ Bark	▪ Tannins	
<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Arn. (Arjun) [107]	▪ Bark	▪ Tannin ▪ Saponin ▪ Flavonoids ▪ Gallic acid	▪ Control glucose metabolizing enzyme activities
<i>Terminalia bellirica</i> (Gaertn.) Roxb. (Bahera) [108, 109]	▪ Fruit ▪ Leaf		▪ Stimulate insulin secretion ▪ Inhibit activities of α -amylase and α -glucosidase
<i>Terminalia chebula</i> Retz. (Haritaki) [110, 111]	▪ Fruit	▪ Tannins ▪ Triterpenoids ▪ Flavonoids ▪ Flavonoids	▪ Regulate the activities of glucose metabolizing enzymes ▪ Regenerate pancreatic β -cells ▪ Inhibit activities of α -glucosidase ▪ Inhibit α -glucosidase activities
<i>Vigna radiata</i> (L.) R. Wilczek (Mug) [112]	▪ Seed		
<i>Vigna unguiculata</i> (L.) Walp. (Barbati) [113]	▪ Seed	▪ Fibers	▪ Activate the insulin signaling cascade
<i>Vitex negundo</i> L. (Nishinda) [114, 115]	▪ Leaf	▪ Idopyranose ▪ Iridoid glucoside	▪ Regenerate β -cells of the pancreas ▪ Enhance pancreatic insulin secretion

<i>Withania somnifera</i> (L.) Dunal (Aswagandha) [116, 117]	▪ Root	▪ Flavonoids	▪ Regulate carbohydrate metabolizing enzyme activity
<i>Xylocarpus moluccensis</i> (Lam.) M. Roem. (Passur) [118]	▪ Leaf		▪ Improve insulin sensitivity
<i>Zea mays</i> L. (Bhootta) [119, 120]	▪ Fruit	▪ Limonoids	▪ Reduce the activity of glucose-6-phosphatase
			▪ Alter the activities of glucose metabolizing enzymes
			▪ Increase glucose uptake by muscle cells
			▪ Inhibit α -glucosidase activities
			▪ Inhibit α -glucosidase activities
			▪ Protect pancreatic β -cells
			▪ Increase insulin secretion
<i>Zingiber officinale</i> Roscoe (Ada) [121, 122]	▪ Kernel	▪ Phenolics	▪ Increase the activities of glycolytic enzymes
		▪ Flavonoids	▪ Enhance Glucose uptake
	▪ Rhizome	▪ Gingerol	

CONCLUSION

Since the advent of human beings, plants continue to be a vital component not only for fostering human health and well-being but also for flourishing the quality of human life via alleviating ailments. Several studies have reported the utility of a number of plants for the cessation and mitigation of many chronic diseases, including DM. So, researchers are pursuing among the native traditional medicinal plants in search of a substitute to antidiabetic drugs owing to the unwanted side-effects, availability, and accessibility of conventional antihyperglycemic drugs. Bangladesh, being a fertile and alluvial land exaggerated with tropical forests and boggy jungles, is an affluent source of a wide variety of medicinal plants applied in the traditional medicinal system for curing disease.

The current review has presented a comprehensive detail of the antidiabetic plants being offered in the historical platform of Bangladeshi medication for management and treatment of DM. The results demonstrated a few pharmacological mechanisms of actions of the medicinal plants to accomplish their therapeutic actions. *In vivo* experiments displayed that inhibition of absorption of glucose in intestinal, an increment in the biosynthesis of liver glycogen or decrement in glycogenolysis, improvement in peripheral glucose absorption etc. are among some antidiabetic mechanistic actions of Bangladeshi natural remedies. Rejuvenation of pancreatic β -cells, inhibiting the decay of islet tissue of pancreas, increase in the restrictive action against α -amylase and α -glucosidase enzyme, a rise of insulin sensitivity or the insulin-like performance of the plant essence is the *in vitro* findings supporting the dexterity of the plants used for DM. It is suggested that the glycemic control properties of the plants are attributed to the bioactive principles like alkaloids, flavonoids terpenoids, glycosides etc. These secondary plant metabolites have already proven their role in preventing and treating many chronic diseases like diabetes. Although, among the clinical experiments, no serious hostile effects were investigated and thus, the natural herbal preparations in gross assumed harmless for the human. However, every plant material is not safe, and so, more investigation of the noxious effect of these herbal plants is needed prior to recommending for human use.

In conclusion, the current review contributes to an insight of the prevailing perception of the efficiency of plant extracts commonly adopted in Bangladesh as a dietary supplement or ancillary therapy for the preclusion and remedy of diabetes. Nonetheless, the evidence found so far is mainly suggestive, and thus, it is crucial to recognize the bioactive component accountable for the antihyperglycemic action with the actual mode of action. More specific and focused research, therefore, will provide a better and broad understanding in this regard. Lastly, this extensive literature review can serve as an efficient mean for the nomination of plants having an intense potential for the unbolting of novel antidiabetic agents.

Conflict of interest

The authors' have no conflict of interest to declare.

REFERENCES

- ADA. Diagnosis and classification of diabetes mellitus. American Diabetes Association. Diabetes Care. 2014; 37(Suppl 1):S81-S90.
- WHO. Global report on diabetes. World Health Organization, 2016.
- Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. PLoS Med. 2006; 3:e442.
- Akter S, Rahman MM, Abe SK, Sultana P. Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. Bull World Health Organ. 2014; 92:204-213.
- Kirpichnikov D, McFarlane SI, Sowers JR. Metformin: an update. Ann Intern Med. 2002; 137:25-33.
- Lorenzati B, Zucco C, Miglietta S, Lamberti F, Bruno G. Oral Hypoglycemic Drugs: Pathophysiological Basis of Their Mechanism of Action Oral Hypoglycemic Drugs: Pathophysiological Basis of Their Mechanism of Action. Pharmaceuticals (Basel). 2010; 3:3005-3020.
- Ghorbani A. Clinical and experimental studies on polyherbal formulations for diabetes: current status and future prospective. J Integr Med 2014; 12:336-345.
- Ghorbani A. Best herbs for managing diabetes: A review of clinical studies. Braz J Pharm Sci 2013; 49:413-422.
- Venkatesh S, Reddy GD, Reddy BM, Ramesh M, Rao AV. Antihyperglycemic activity of *Caralluma attenuata*. Fitoterapia. 2003; 74:274-279.
- Dixie G, Imam SA, Hussain MJ. Medicinal plant marketing in Bangladesh. Dhaka: Intercooperation Bangladesh. 2003, 8-22.
- Yusuf M, Begum J, Hoque MN, Choudhury JU. Medicinal plants of Bangladesh-Revised and Enlarged. Bangladesh Centre for Scientific and Industrial Research, Chittagong, Bangladesh, 2009.
- Rahman A. Ethno-botanical Survey of Traditional Medicine Practice for the Treatment of Cough, Diabetes, Diarrhea, Dysentery and Fever of Santals at Abdullahpur Village under Akkelpur Upazilla of Joypurhat District, Bangladesh. Biomed Biotechnol. 2013; 1:27-30.
- Rahman M, Mishuk A, Halder S, Rouf ASS. A Survey on Traditional Medicinal Plants used for the Treatment of Diabetes in Urban Areas of Dhaka and Khulna, Bangladesh. Glob J Med Res 2013; 13.
- Rahman AM. Ethno-botanical Survey of Anti-Diabetic Medicinal Plants Used by the Santal Tribe of Joypurhat District, Bangladesh. Int J Res Pharm Biosci 2015; 2:19-26.
- Rahman AM, Keya MA. Traditional Medicinal Plants Used by local People at the Village Sabgram under Sadar Upazila of Bogra District, Bangladesh. Res Plant Sci. 2015; 3:31-37.
- Ray D, Sharatchandra K, Thokchom I. Antipyretic, antidiarrhoeal, hypoglycaemic and hepatoprotective activities of ethyl acetate extract of *Acacia catechu* Willd. in albino rats. Indian J Pharm 2006; 38:408.
- Gao H, Huang YN, Gao B, Li P, Inagaki C, Kawabata J. Inhibitory effect on α -glucosidase by *Adhatoda vasica* Nees. Food Chem. 2008; 108:965-972.
- Kamalakkannan N, Prince P. Antidiabetic and anti-oxidant activity of *Aegle marmelos* extract in streptozotocin-induced diabetic rats. Pharm Biol. 2004; 42:125-130.
- Gandhi GR, Ignacimuthu S, Paulraj MG. Hypoglycemic and beta-cells regenerative effects of *Aegle marmelos* (L.) Corr. bark extract in streptozotocin-induced diabetic rats. Food Chem Toxicol. 2012; 50:1667-1674.
- Kumari K, Mathew BC, Augusti KT. Antidiabetic and hypolipidemic effects of S-methyl cysteine sulfoxide isolated from *Allium cepa* Linn. Indian J Biochem Biophys 1995; 32:49-54.

21. Jung JY, Lim Y, Moon MS, Kim JY, Kwon O. Onion peel extracts ameliorate hyperglycemia and insulin resistance in high fat diet/streptozotocin-induced diabetic rats. *Nutr Metab (Lond)*. 2011; 8:18.
22. Sheela CG, Augusti KT. Antidiabetic effects of S-allyl cysteine sulphoxide isolated from garlic *Allium sativum* Linn. *Indian J Exp Biol* 1992; 30:523-526.
23. Eidi A, Eidi M, Esmaeili E. Antidiabetic effect of garlic (*Allium sativum* L.) in normal and streptozotocin-induced diabetic rats. *Phytomedicine*. 2006; 13:624-629.
24. Rahman MM, Hossain MA, Siddique SA, Biplab KP, Uddin MH. Antihyperglycemic, antioxidant and cytotoxic activities of *Alocasia macrorrhizos* (Linn.) rhizomes extract. *Turk J Biol* 2012; 36:574-579.
25. Rajasekaran S, Sivagnanam K, Ravi K, Subramanian S. Hypoglycemic effect of Aloe vera gel on streptozotocin-induced diabetes in experimental rats. *J Med Food* 2004; 7:61-66.
26. Ajabnoor MA. Effect of aloe on blood glucose levels in normal and alloxan diabetic mice. *J Ethnopharmacol* 1990; 28:215-220.
27. Ashok Kumar BS, Lakshman K, Nandeesh R, Arun Kumar PA, Manoj B, Kumar V, *et al.* *In vitro* alpha-amylase inhibition and *in vivo* antioxidant potential of *Amaranthus spinosus* in alloxan-induced oxidative stress in diabetic rats. *Saudi J Biol Sci* 2011; 18:1-5.
28. Mishra SB, Verma A, Mukerjee A, Vijayakumar M. *Amaranthus spinosus* L.(Amaranthaceae) leaf extract attenuates streptozotocin-nicotinamide induced diabetes and oxidative stress in albino rats: A histopathological analysis. *Asian Pac J Trop Biomed* 2012; 2:S1647-S1652.
29. Ashok Kumar BS, Lakshman K, Jayaveea KN, Sheshadri Shekar D, Saleemulla K, Thippeswamy BS, *et al.* Antidiabetic, antihyperlipidemic and antioxidant activities of methanolic extract of *Amaranthus viridis* Linn in alloxan induced diabetic rats. *Exp Toxicol Pathol*. 2012; 64:75-79.
30. Uddin S, Islam MM, Hassan MM, Bhowmik A, Rokeya B. *Amaranthus viridis* modulates anti-hyperglycemic pathways in hemi-diaphragm and improves glycogenesis liver function in rats. *J Pharmacogn Phytother* 2016; 8:173-181.
31. Reddy AH, Jamuna JB, Salimath P, Aradhya S. Elephant foot yam ameliorates activities of intestinal and renal disaccharidases in streptozotocin induced diabetic rats. *South Asian J Exp Biol* 2012; 2:141-148.
32. Xie W, Wang W, Su H, Xing D, Pan Y, Du L. Effect of ethanolic extracts of *Ananas comosus* L. leaves on insulin sensitivity in rats and HepG2. *Comp Biochem Physiol C Toxicol Pharmacol*. 2006; 143:429-435.
33. Kalpana M, Prasath G, Subramanian S. Studies on the antidiabetic activity of *Ananas comosus* leaves in STZ induced diabetic rats. *Der Pharmacia Lett*. 2014; 6:190-198.
34. Subramanian R, Asmawi MZ, Sadikun A. *In vitro* alpha-glucosidase and alpha-amylase enzyme inhibitory effects of *Andrographis paniculata* extract and andrographolide. *Acta Biochim Pol*. 2008; 55:391-398.
35. Zhang X-F, Tan B-h. Anti-diabetic property of ethanolic extract of *Andrographis paniculata* in streptozotocin-diabetic rats. *Acta Pharmacol Sin*. 2000; 21:1157-1164.
36. Amudhan M, Begum V. Alpha-glucosidase inhibitory and hypoglycemic activities of *Areca catechu* extract. *Pharmacogn Mag*. 2008; 4:223.
37. Hannan JM, Ali L, Khaleque J, Akhter M, Flatt PR, Abdel-Wahab YH. Antihyperglycaemic activity of *Asparagus racemosus* roots is partly mediated by inhibition of carbohydrate digestion and absorption, and enhancement of cellular insulin action. *Br J Nutr* 2012; 107:1316-1323.
38. Satyanarayana K, Sravanthi K, Shaker IA, Ponnulakshmi R. Molecular approach to identify antidiabetic potential of *Azadirachta indica*. *J Ayurveda Integr Med* 2015; 6:165-174.
39. Khosla P, Bhanwra S, Singh J, Seth S, Srivastava RK. A study of hypoglycaemic effects of *Azadirachta indica* (Neem) in normal and alloxan diabetic rabbits. *Indian J Physiol Pharmacol* 2000; 44:69-74.
40. Kochhar A, Sharma N, Sachdeva R. Effect of supplementation of *Tulsi* (*Ocimum sanctum*) and *Neem* (*Azadirachta indica*) leaf powder on diabetic symptoms, anthropometric parameters and blood pressure of non insulin dependent male diabetics. *Ethno-Med*. 2009; 3:5-9.
41. Khatib N, Patil P. Evaluation of hypoglycemic activity of *Barringtonia acutangula* fruit extracts in streptozotocin induced hyperglycemic wistar rats. *J Cell Tissue Res* 2011; 11:2573.
42. Babre NP, Debnath S, Manjunath S, Reddy VM, Murlidharan P, Manoj G. Antidiabetic effect of hydroalcoholic extract of *Barringtonia acutangula* Linn. Root on streptozotocin induced diabetic rats. *Int J Pharm Sci Nanotechnol* 2010; 3:1158-1164.
43. Thavamani BS, Subburaj V. *In vitro* Studies on *Basella rubra* Different Extracts as Inhibitors of Key Enzymes Linked to Diabetes Mellitus. *Pharmacognosy J* 2017; 9.
44. Pari L, Satheesh MA. Antidiabetic activity of *Boerhaavia diffusa* L.: effect on hepatic key enzymes in experimental diabetes. *J Ethnopharmacol* 2004; 91:109-113.
45. Mir MA, Mir BA, Bisht A, Rao Z, Singh D. Antidiabetic properties and metal analysis of *Bombax ceiba* flower extracts. *Glob J Addict Rehabil Med* 2017; 1.
46. Somani R, Kasture S, Singhai AK. Antidiabetic potential of *Butea monosperma* in rats. *Fitoterapia*. 2006; 77:86-90.
47. Panda S, Jafri M, Kar A, Meheta BK. Thyroid inhibitory, antiperoxidative and hypoglycemic effects of stigmaterol isolated from *Butea monosperma*. *Fitoterapia*. 2009; 80:123-126.
48. Wang Y, Yang Z, Wei X. Sugar compositions, alpha-glucosidase inhibitory and amylase inhibitory activities of polysaccharides from leaves and flowers of *Camellia sinensis* obtained by different extraction methods. *Int J Biol Macromol* 2010; 47:534-539.
49. Veeramani C, Pushpavalli G, Pugalendi KV. Antihyperglycaemic effect of *Cardiospermum halicacabum* Linn. leaf extract on STZ-induced diabetic rats. *J Appl Biomed* 2008; 6.
50. Juarez-Rojop IE, Diaz-Zagoya JC, Ble-Castillo JL, Miranda-Osorio PH, Castell-Rodriguez AE, Tovilla-Zarate CA, *et al.* Hypoglycemic effect of *Carica papaya* leaves in streptozotocin-induced diabetic rats. *BMC Complement Altern Med*. 2012; 12:236.
51. Singh SN, Vats P, Suri S, Shyam R, Kumria MM, Ranganathan S, *et al.* Effect of an antidiabetic extract of *Catharanthus roseus* on enzymic activities in streptozotocin induced diabetic rats. *J Ethnopharmacol* 2001; 76:269-277.
52. Rasineni K, Bellamkonda R, Singareddy SR, Desireddy S. Antihyperglycemic activity of *Catharanthus roseus* leaf powder in streptozotocin-induced diabetic rats. *Pharmacognosy Res*. 2010; 2:195-201.
53. Jayanthi M, Sowbala N, Rajalakshmi G, Kanagavalli U, Sivakumar V. Study of antihyperglycemic effect of *Catharanthus roseus* in alloxan induced diabetic rats. *Int J Pharm Pharm Sci* 2010; 2:114-116.
54. Supkamonseni N, Thinkratok A, Meksuriyen D, Srisawat R. Hypolipidemic and hypoglycemic effects of *Centella asiatica* (L.) extract *in vitro* and *in vivo*. *Indian J Exp Biol* 2014; 52:965-971.
55. Kabir AU, Samad MB, D'Costa NM, Akhter F, Ahmed A, Hannan JM. Antihyperglycemic activity of *Centella asiatica* is partly mediated by carbohydrate inhibition and glucose-fiber binding. *BMC Complement Altern Med*. 2014; 14:31.
56. Kumanan R, Manimaran S, Saleemulla K, Dhanabal S, Nanjan M. Screening of bark of *Cinnamomum tamala* (Lauraceae) by using α -amylase inhibition assay for anti-diabetic activity. *Int J Pharm Biome Res* 2010; 1:69-72.
57. Gupta R, Agnihotri PK, Johri S, Saxena AM. Hypoglycaemic activity of ethanol extract of *Cinnamomum tamala* leaves in normal and streptozotocin diabetic rats. *Iran J Pharmacol Ther* 2009; 8:17-21.
58. Hassan SA, Barthwal R, Nair MS, Haque SS. Aqueous Bark Extract of *Cinnamomum Zeylanicum*: A Potential Therapeutic Agent for Streptozotocin- Induced Type 1 Diabetes Mellitus (T1DM) Rats. *Trop J Pharm Res* 2012; 11:429-435.
59. Daisy P, Rajathi M. Hypoglycemic effects of *Clitoria ternatea* Linn. (Fabaceae) in alloxan-induced diabetes in rats. *Trop J Pharm Res* 2009; 8:393-398.
60. Verma PR, Itankar PR, Arora SK. Evaluation of antidiabetic antihyperlipidemic and pancreatic regeneration, potential of aerial parts of *Clitoria ternatea*. *Rev Bras Farmacogn*. 2013; 23:819-829.
61. Attanayake A, Jayatilaka K, Pathirana C, Mudduwa L. Antihyperglycemic activity of *Coccinia grandis* (L.) Voigt in streptozotocin induced diabetic rats. *Indian J Tradit Knowl* 2015; 14:376-381.
62. Mohammed SI, Chopda MZ, Patil RH, Vishwakarma KS, Maheshwari VL. *In vivo* antidiabetic and antioxidant activities of *Coccinia grandis* leaf extract against streptozotocin induced diabetes in experimental rats. *Asian Pac J Trop Dis* 2016; 6:298-304.
63. Lekshmi PC, Arimboor R, Indulekha PS, Menon AN. Turmeric (*Curcuma longa* L.) volatile oil inhibits key enzymes linked to type 2 diabetes. *Int J Food Sci Nutr* 2012; 63:832-834.
64. Krishna Murthy B, Nammi S, Kota MK, Krishna Rao RV, Koteswara Rao N, Annapurna A. Evaluation of hypoglycemic and antihyperglycemic effects of *Datura metel* (Linn.) seeds in normal and alloxan-induced diabetic rats. *J Ethnopharmacol* 2004; 91:95-98.
65. Kumar S, Kumar V, Prakash O. Enzymes inhibition and antidiabetic effect of isolated constituents from *Dillenia indica*. *Biomed Res Int*. 2013; 2013:382063.
66. Gayathri M, Kannabiran K. Antidiabetic and ameliorative potential of *Ficus bengalensis* bark extract in streptozotocin induced diabetic rats. *Indian J Clin Biochem*. 2008; 23:394-400.

67. Anitha T, Balakumar C, Ilango K, Jose CB, Vetrivel D. Antidiabetic activity of the aqueous extracts of *Foeniculum vulgare* on streptozotocin-induced diabetic rats. *Int J Adv Pharm Biol Chem* 2014; 3:487-494.
68. Venkatesh S, Thilagavathi J, Shyam Sundar D. Anti-diabetic activity of flowers of *Hibiscus rosasinensis*. *Fitoterapia*. 2008; 79:79-81.
69. El-Sawi N, Gad MH, Al-Seeni MN, Younes S, Soad E-ME-Ga, Ali S. Evaluation of Antidiabetic Activity of *Ipomoea Aquatica* Fractions in Streptozotocin Induced Diabetic in Male Rat Model. *Sohag J Sci* 2017; 2:9-17.
70. El-Baz F, Aly H, Abd-Alla H, Saad S. Bioactive flavonoid glycosides and antidiabetic activity of *Jatropha curcas* on streptozotocin-induced diabetic rats. *Int J Pharm Sci Rev Res* 2014; 29:143-156.
71. Patil SB, Dongare VR, Kulkarni CR, Joglekar MM, Arvindekar AU. Antidiabetic activity of *Kalanchoe pinnata* in streptozotocin-induced diabetic rats by glucose independent insulin secretagogue action. *Pharm Biol*. 2013; 51:1411-1418.
72. Yuliani T, Dewijanti ID, Banjarnahor SDS. Antidiabetic Activity of Ethanolic Extract of *Kalanchoe Pinnata* Leaves in Alloxan Induced Hyperglycaemic Rats. *Indonesian J Pharm* 2016; 27:139-144.
73. Matthew S, Singh D, Jaiswal S, M.Kumar.B.Jayakar, Bhowmik D. Antidiabetic Activity of *Kalanchoe Pinnata* (Lam.) Pers in alloxan Induced Diabetic Rats. *J Chem Pharm Sci* 2013; 6:1-7.
74. Bhattacharya S, Das B. Anti-diabetic activity of *Lagenaria siceraria* pulp and seed extract in normal and alloxan-induced diabetic rats. *Int J Pharm Sci Res* 2012; 3:3362.
75. Saha BK, Bhuiyan MNH, Mazumder K, Haque KF. Hypoglycemic activity of *Lagerstroemia speciosa* L. extract on streptozotocin-induced diabetic rat: Underlying mechanism of action. *Bangladesh J Pharmacol* 2009; 4:79-83.
76. Imam H, Mahbub NU, Khan MF, Hana HK, Sarker MM. Alpha amylase enzyme inhibitory and anti-inflammatory effect of *Lawsonia inermis*. *Pak J Biol Sci* 2013; 16:1796-1800.
77. Ghule AE, Jadhav SS, Bodhankar SL. Effect of ethanolic extract of seeds of *Linum usitatissimum* (Linn.) in hyperglycaemia associated ROS production in PBMCs and pancreatic tissue of alloxan induced diabetic rats. *Asian Pac J Trop Dis* 2012; 2:405-410.
78. Pimple B, Kadam P, Patil M. Antidiabetic and antihyperlipidemic activity of *Luffa acutangula* fruit extracts in streptozotocin induced niddm rats. *Asian J Pharm Clin Res* 2011; 4:156-163p.
79. Chaurasia S, Saxena R, Chaurasia I, Shrivastav R. Antidiabetic activity of *Luffa aegyptica* (Mill) in alloxan induced diabetic rats. *J Chem Pharm Res* 2011; 3:522-525.
80. Ganapaty S, Nandeesh R, Veerapur V, Thippeswamy B, Shivasharan D. In vivo and In vitro Anti-Diabetic Effects of *Madhuca indica* Roxb., in Alloxan-Induced Diabetic rats. *Int J Adv Pharm Biol Chem* 2013; 2:282-290.
81. Ganogpichayagrai A, Palanuvej C, Ruangrunsi N. Antidiabetic and anticancer activities of *Mangifera indica* cv. Okrong leaves. *J Adv Pharm Technol Res* 2017; 8:19-24.
82. Kazeem MI, Ogungbe SM, Saibu GM, Aboyade OM. In vitro study on the hypoglycemic potential of *Nicotiana tabacum* leaf extracts. *Bangladesh J Pharmacol* 2014; 9:140-145.
83. Benhaddou-Andaloussi A, Martineau LC, Spoor D, Vuong T, Leduc C, Joly E, et al. Antidiabetic activity of *Nigella sativa*. Seed extract in cultured pancreatic β -cells, skeletal muscle cells, and adipocytes. *Pharm Biol*. 2008; 46:96-104.
84. Fallah Huseini H, Amini M, Mohtashami R, Ghamarchehre ME, Sadeqhi Z, Kianbakht S, et al. Blood pressure lowering effect of *Nigella sativa* L. seed oil in healthy volunteers: a randomized, double-blind, placebo-controlled clinical trial. *Phytother Res*. 2013; 27:1849-1853.
85. Bamosa AO, Kaatabi H, Lebdaa FM, Elq AM, Al-Sultanb A. Effect of *Nigella sativa* seeds on the glycemic control of patients with type 2 diabetes mellitus. *Indian J Physiol Pharmacol* 2010; 54:344-354.
86. Huang YN, Zhao YL, Gao XL, Zhao ZF, Jing Z, Zeng WC, et al. Intestinal alpha-glucosidase inhibitory activity and toxicological evaluation of *Nymphaea stellata* flowers extract. *J Ethnopharmacol* 2010; 131:306-312.
87. Rajagopal K, Sasikala K. Antidiabetic activity of hydro-ethanolic extracts of *Nymphaea Stellata* flowers in normal and alloxan induced diabetic rats. *Afr J Pharm Pharmacol* 2008; 2:173-178.
88. Hannan J, Mareneh L, Ali L, Rokeya B, Flatt P, Abdel-Wahab Y. *Ocimum sanctum* leaf extracts stimulate insulin secretion from perfused pancreas, isolated islets and clonal pancreatic β -cells. *J Endocrinol* 2006; 189:127-136.
89. Vats V, Yadav SP, Grover JK. Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozotocin-induced alterations in glycogen content and carbohydrate metabolism in rats. *J Ethnopharmacol* 2004; 90:155-160.
90. Boue SM, Daigle KW, Chen MH, Cao H, Heiman ML. Antidiabetic Potential of Purple and Red Rice (*Oryza sativa* L.) Bran Extracts. *J Agric Food Chem* 2016; 64:5345-5353.
91. Kumari S, Wanjari M, Kumar P, Palani S. Antidiabetic activity of *Pandanus fascicularis* Lamk-aerial roots in alloxan-induced hyperglycemic rats. *Int J Nutr Pharmacol Neurol Dis* 2012; 2:105.
92. Sultana Z, Jami SI, Ali E, Begum M, Haque M. Investigation of Antidiabetic Effect of Ethanolic Extract of *Phyllanthus emblica* Linn. Fruits in Experimental Animal Models. *Pharmacol Pharm*. 2014; 5:11.
93. Santhakumari P, Prakasam A, Pugalendi KV. Antihyperglycemic activity of Piper betle leaf on streptozotocin-induced diabetic rats. *J Med Food* 2006; 9:108-112.
94. Sivashanmugam AT, Chatterjee TK. In vitro and in vivo antidiabetic activity of *Polyalthia longifolia* (Sonner.) Thw. leaves. *Orient Pharm Exp Med*. 2013; 13:289-300.
95. Khalil EA. Antidiabetic effect of an aqueous extract of Pomegranate (*Punica granatum* L.) peels in normal and alloxan diabetic rats. *Egypt J Hosp Med* 2004; 16:92-99.
96. Mishra S, Vijayakumar M. Anti-hyperglycemic and antioxidant effect of *Saraca asoca* (Roxb. De Wilde) flowers in Streptozotocin-Nicotinamide induced diabetic rats: A therapeutic study. *J Bioanal Biomed S* 2014; 12.
97. Kumar S, Narwal S, Kumar D, Singh G, Narwal S, Arya R. Evaluation of antihyperglycemic and antioxidant activities of *Saraca asoca* (Roxb.) De Wild leaves in streptozotocin induced diabetic mice. *Asian Pac J Trop Dis* 2012; 2:170-176.
98. Kwon YI, Apostolidis E, Shetty K. In vitro studies of eggplant (*Solanum melongena*) phenolics as inhibitors of key enzymes relevant for type 2 diabetes and hypertension. *Bioresour Technol*. 2008; 99:2981-2988.
99. Gomathi V, Jayakar B, Kothai R, Ramakrishnan G. Antidiabetic activity of leaves of *Spinacia oleracea* Linn. *J Chem Pharm Res* 2010; 2:266-274.
100. Attanayake AP, Jayatilaka KAPW, Pathirana C, Mudduwa LKB. Antihyperglycaemic, antihyperlipidaemic and β cell regenerative effects of *Spondias pinnata* (Linn. f.) Kurz. bark extract on streptozotocin induced diabetic rats. *Eur J Integr Med* 2014; 6:588-596.
101. Shinde J, Taldone T, Barletta M, Kunaparaju N, Hu B, Kumar S, et al. Alpha-glucosidase inhibitory activity of *Syzygium cumini* (Linn.) Skeels seed kernel in vitro and in Goto-Kakizaki (GK) rats. *Carbohydr Res*. 2008; 343:1278-1281.
102. Wang W, Xu H, Chen H, Tai K, Liu F, Gao Y. In vitro antioxidant, anti-diabetic and antilipemic potentials of quercetagenin extracted from marigold (*Tagetes erecta* L.) inflorescence residues. *J Food Sci Technol* 2016; 53:2614-2624.
103. Maiti R, Jana D, Das UK, Ghosh D. Antidiabetic effect of aqueous extract of seed of *Tamarindus indica* in streptozotocin-induced diabetic rats. *J Ethnopharmacol* 2004; 92:85-91.
104. Hamidreza H, Heidari Z, Shahraki M, Moudi B. A stereological study of effects of aqueous extract of *Tamarindus indica* seeds on pancreatic islets in streptozotocin-induced diabetic rats. *Pak J Pharm Sci* 2010; 23:427-434.
105. Parvin A, Alam MM, Haque MA, Bhowmik A, Ali L, Rokeya B. Study of the Hypoglycemic Effect of *Tamarindus indica* Linn. Seeds on Non-Diabetic and Diabetic Model Rats. *Br J Pharm Res* 2013; 3:1094-1105.
106. Varma SB, Jaybhaye DL. Antihyperglycemic activity of *Tectona grandis* Linn. bark extract on alloxan induced diabetes in rats. *Int J Ayurveda Res* 2010; 1:163-166.
107. Ragavan B, Krishnakumari S. Antidiabetic effect of *T. arjuna* bark extract in alloxan induced diabetic rats. *Indian J Clin Biochem* 2006; 21:123-128.
108. Kasabri V, Flatt PR, Abdel-Wahab YH. *Terminalia bellirica* stimulates the secretion and action of insulin and inhibits starch digestion and protein glycation in vitro. *Br J Nutr* 2010; 103:212-217.
109. Manila T, Mary Shoba Das C, Gayathri Devi S. In vitro anti-diabetic properties of the fruits and leaves of *Terminalia bellirica*. *J Nat Sci Res* 2012; 2:54-62.
110. Kumar GPS, Arulselvan P, Kumar DS, Subramanian SP. Anti-diabetic activity of fruits of *Terminalia chebula* on streptozotocin induced diabetic rats. *J Health Sci* 2006; 52:283-291.
111. Gao H, Huang Y-N, Xu P-Y, Kawabata J. Inhibitory effect on α -glucosidase by the fruits of *Terminalia chebula* Retz. *Food Chem*. 2007; 105:628-634.
112. Yao Y, Yang X, Tian J, Liu C, Cheng X, Ren G. Antioxidant and antidiabetic activities of black mung bean (*Vigna radiata* L.). *J Agric Food Chem* 2013; 61:8104-8109.
113. Barnes M, Uruakpa F, Udenigwe C. Influence of cowpea (*Vigna unguiculata*) peptides on insulin resistance. *J Nutr Health Food Sci* 2015; 3:1-3.
114. Manikandan R, Thiagarajan R, Beulaja S, Sivakumar MR, Meiyalagan V, Sundaram R, et al. 1, 2 di-substituted idopyranose from *Vitex negundo* L.

- protects against streptozotocin-induced diabetes by inhibiting nuclear factor-kappa B and inducible nitric oxide synthase expression. *Microsc Res Tech.* 2011; 74:301-307.
115. Sundaram R, Naresh R, Ranadevan R, Shanthi P, Sachdanandam P. Effect of iridoid glucoside on streptozotocin induced diabetic rats and its role in regulating carbohydrate metabolic enzymes. *Eur J Pharmacol* 2012; 674:460-467.
 116. Anwer T, Sharma M, Pillai KK, Iqbal M. Effect of *Withania somnifera* on insulin sensitivity in non-insulin-dependent diabetes mellitus rats. *Basic Clin Pharmacol Toxicol.* 2008; 102:498-503.
 117. Udayakumar R, Kasthuriangan S, Mariashibu TS, Rajesh M, Anbazhagan VR, Kim SC, *et al.* Hypoglycaemic and hypolipidaemic effects of *Withania somnifera* root and leaf extracts on alloxan-induced diabetic rats. *Int J Mol Sci* 2009; 10:2367-2382.
 118. Srivastava AK, Tiwari P, Srivastava SP, Srivastava R, Mishra A, Rahuja N, *et al.* Antihyperglycaemic and antidyslipidemic activities in ethyl acetate fraction of fruits of marine mangrove *Xylocarpus moluccensis*. *Int J Pharm Pharm Sci* 2014; 6:809-826.
 119. Nile SH, Park SW. Antioxidant, alpha-glucosidase and xanthine oxidase inhibitory activity of bioactive compounds from maize (*Zea mays* L.). *Chem Biol Drug Des.* 2014; 83:119-125.
 120. Huang B, Wang Z, Park JH, Ryu OH, Choi MK, Lee JY, *et al.* Anti-diabetic effect of purple corn extract on C57BL/KsJ db/db mice. *Nutr Res Pract.* 2015; 9:22-29.
 121. Abdulrazaq NB, Cho MM, Win NN, Zaman R, Rahman MT. Beneficial effects of ginger (*Zingiber officinale*) on carbohydrate metabolism in streptozotocin-induced diabetic rats. *Br J Nutr* 2012; 108:1194-1201.
 122. Li Y, Tran VH, Duke CC, Roufogalis BD. Gingerols of *Zingiber officinale* enhance glucose uptake by increasing cell surface GLUT4 in cultured L6 myotubes. *Planta Med.* 2012; 78:1549-1555.

HOW TO CITE THIS ARTICLE

Hasan T, Sultana M. Antidiabetic Potency of Bangladeshi Medicinal Plants. *J Ayu Herb Med* 2018;4(1):35-42.