



## Review Article

ISSN: 2454-5023  
J. Ayu. Herb. Med.  
2018; 4(1): 43-47  
© 2018, All rights reserved  
www.ayurvedjournal.com  
Received: 03-02-2018  
Accepted: 03-04-2018

## In-silico studies in herbal drugs: A review

Dr. Monisha.M<sup>1</sup>, Dr. Abhaya Kumar Mishra<sup>2</sup>, Dr. Ramesh N V<sup>3</sup>, Dr. K Unnikrishna Pillai<sup>4</sup>, Dr. Vineeth P K<sup>5</sup>

<sup>1</sup> PG Scholar, Department of Rasashastra and Bhaishajya Kalpana (Medicinal chemistry and pharmacy), Amrita School of Ayurveda, Amritapuri; Amrita Viswa Vidyapeetham; India

<sup>2</sup> Professor, Department of Rasashastra and Bhaishajya Kalpana (Medicinal chemistry and pharmacy), Amrita School of Ayurveda, Amritapuri; Amrita Viswa Vidyapeetham; India

<sup>3</sup> Professor and HOD, Department of Rasashastra and Bhaishajya Kalpana (Medicinal chemistry and pharmacy), Amrita School of Ayurveda, Amritapuri; Amrita Viswa Vidyapeetham; India

<sup>4</sup> Professor, Department of Rasashastra and Bhaishajya Kalpana (Medicinal chemistry and pharmacy), Amrita School of Ayurveda, Amritapuri; Amrita Viswa Vidyapeetham; India

<sup>5</sup> Assistant Professor, Department of Rasashastra and Bhaishajya Kalpana, (Medicinal chemistry and pharmacy), Amrita School of Ayurveda, Amritapuri; Amrita Viswa Vidyapeetham; India

### ABSTRACT

Bioinformatics is the latest technology in the research field. In-silico studies are done to identify the exact target of the drug. Which finds a drug for the particular binding site and final stage animal testing can be done for obtaining a conform result. Reviews of in-silico studies in herbal drugs were retrieved through the use of PubMed. Specific software on a computer allows researchers to analyse enormous data without actually conducting a large number of experiments. It helps to give the existing information to model disease pathway and identifies precise targets of the selected drugs. Modern instrumental techniques like XRD, XRF, ICP-MS etc. help in quantitative and qualitative estimation of metals and minerals and structural analysis of compound drugs. A later stage in vivo and in vitro studies can be done for obtaining the confirmatory result.

**Keywords:** *In-silico*, Herbal drugs, Bioinformatics.

### INTRODUCTION

Using a set of complex pharmaceutical combinations in treatment is one of the most important aspects of therapeutics. So, to evaluate the proper therapeutic value or the clinical efficacy of the multicomponent/polyherbal formulations needs modern technology. Bioinformatics plays an important role in medical science by identifying the exact cell, genes, proteins being affected and to screen the active protein among them. Biology has been transformed into an information science where molecular processes can be addressed computationally. In the research field, *in-vivo* and *in-vitro* studies have an important role before the clinical trials of a new medicine. So, to confirm the target cell of the particular medicine the bioinformatics plays a vital role before going to *in-vivo* studies.

### Methodology through Bioinformatics

1. The test sample is prepared according to the reference
2. The test sample was analysed for structural changes and ionic forms using analytical techniques like XRD, XRF, NMR, IR at a recognized laboratory.
3. The analysed data was used for the bioinformatics software e.g. Linux for the analysis.
4. Based on the structures identified by the above analytical techniques, the computer program generates a structural formula and utilize this structure to assess the protein linkage of the molecule to various proteins in the DNA of human body with various permutations and combinations.
5. The frequency of adhesion of each of the protein linkages is calculated-lesser the frequency faster will be the effect; higher frequency effect was slower of the drug.
6. Like this, we can consider the protein targeted cell.

### *In-silico* Studies in herbal drugs

Post-traumatic stress

An herbal formulation *Free and Easy Wanderer* used as antioxidation therapy in post-traumatic stress

\*Corresponding author:

Dr. Monisha M

PG Scholar, Department of Rasashastra and Bhaishajya Kalpana (Medicinal chemistry and pharmacy), Amrita School of Ayurveda, Amritapuri; Amrita Viswa Vidyapeetham; India  
Email: m.monishanair[at]gmail.com

disorder. Ten compounds selected from the extract counteracted H<sub>2</sub>O<sub>2</sub>-induced oxidative stress [1].

#### Liver disorder

The study of Drug *Radix salviae miltiorrhizae* (Danshen) in liver diseases shows Danshensu exhibits the largest number of hepatoprotective target connections among the six active compounds [2].

#### Cardiovascular diseases

[3] A study on Cardiovascular diseases including thrombosis, which is induced by platelet aggregation. By the usage of 38 compounds have significant potential for development of anti-thrombosis drugs ranked with a DockScore above 70.

[4] One more study shows a positive result of the Danshen drug in cardiovascular and cerebrovascular disease.

[5] Another study on ischemic heart disease treated by Chinese herbal formula *Huo Luo Xiao Ling Dan*. This predicts the molecular interactions between the major active components of apoptosis. [6] Salvianolic acid B from *Salviae miltiorrhizae*, in the treatment of cardiovascular diseases in China.

#### Hypertension

[7] A study done for reducing hypertension by CYP11B2 inhibitor provided a new approach to discover novel CYP11B2 inhibitors from Traditional Chinese Medicine.

[8] In this study Theobroma cacao bean alcoholic extract improves cell viability and reduce IL-6 and sVCAM-1 in endothelial cells induced by plasma (pre-eclamptic patients).

#### Infection & inflammation

[9] Commonly used in anti-infection treatments in Chinese medicine includes the synergistic effects combination of drugs with oxacillin or gentamicin. [10] One more study on Traditional Chinese herbs have a considerable number of drug-like natural products and predicted activities to the targets of approved drugs and usage of these herbs as a chemical resource library for discovery of anti-infective and anti-inflammatory drugs. [11] Another study on extracts from *Ziziphus spinar-christi* for their anti-inflammatory effects.

#### Malaria

[12] Artemisinin is a promising new antimalarial agent.

#### [13] Chikungunya

This study focused on the druggability Nilavembu Kudineer and Swasthya Raksha Amruta Peya to heal virus infection of chikungunya.

#### Tetanus

[14] Phytoligands selected based on literature and pharmacophoric studies. These leads can be used as potential inhibitors against tetanolsin.

#### Myasthenia gravis

[15] Myasthenia gravis treated with Buzhongyiqi decoction shows key active components decoction used to discovered five new AChE inhibitors.

#### Cholesterol

[16] Acyl- coenzyme and Ten compounds finally obtained with predicted inhibitory activities toward ACAT-2.

#### Alopecia.

[17] This present study was aimed to identify prostaglandin D2 synthase inhibitors which is discovered as for treatment of androgenic alopecia.

#### Insomnia

[18] *Kadsura longipedunculata* used for the treatment of insomnia because of its sedative and hypnotic effects.

#### Hemiplegia

[19] Huatuo reconstruction pill treats for hemiplegia and post-operation of brain stroke. [20] This study Identified Amygdalin and paeoniflorin as key active constituents of Buchang Naoxintong in cerebral ischemia..

#### Diabetics mellitus

[21] In the *In-silico* analysis 20 enriched targets predicted for the constituents of *Cassia auriculata* involved in insulin signaling pathways. *Cassia auriculata* enhances uptake of glucose and expression of glucose transporters. [22] Again a study on Plant *Grewia hirsuta* for diabetes. [23] A study to discover novel anti-diabetic drugs. [24] Anti-diabetic drugs have iridoid and secoiridoid glycosides as active chemical components, study shows the inhibition of glycogen phosphorylase- $\alpha$  activity is a common target for iridoids and secoiridoids to elicit anti-diabetic effects. [25] Another study deals with isolated compounds of herb *Pueraria lobata* in type II diabetes mellitus. Compound lobatflavate and 3S,4R-tuberosin shows remarkable  $\alpha$ -glucosidase inhibitory activity.

[26] This study in compounds from *Stephania Hernandifolia* have been confirmed plants against TB and cancer, diabetics.

#### Iron deficiency

[27] A study of piperine in iron deficiency suggests that conjugating piperine with iron slow down the metabolism of iron, thereby piperine probably enhances the bioavailability of iron.

#### Asthma

[28] A study on piperine and its derivatives to detect Lipoxygenase inhibitory activity shows higher significance for medicinal applications.

[29] Another study shows molecular simulation indicates that the MOL376 incough and Asthma.

#### Alzheimer's disease

[30] This study assessed inhibitory action against AChE and BChE and metal-chelating capacity of *Perovskia atriplicifolia* and *Salvia glutinosa*. [31] A study on Anti-Alzheimer's Disease Mechanisms of Icariin. [32] One more study to analyze Anticholinesterase and antioxidant activities of *Terminalia chebula* fruit.

#### Cancer

[33] Scopoletin is constituent in *Artemisia annua* L. Scopoletin might serve as the lead compound for drug development. [34] A study on herbal lead compounds in Prostate Cancer. [35] Another study on Danshen for its anti-cancer effects. [36] Roots of *Rheum undulatum* has having components that are anthraquinone and stilbene derivatives, such as emodin, aloe-emodin, resveratrol, rhaponticin, and isorhapontin and demonstrate sEH inhibitory, antibacterial, antioxidant, anticancer, and anti-inflammatory activity. [37] *Psoralea corylifolia* plant is used for its anti-tumor effects. Major

components in seed is psoralidin. [38] This study Discover novel lead for non-small cell lung cancer. This study suggests the triptolide in Cancer treatment. [39] Another study on liver tumor treated by Fuzheng Yiliu decoction. 11 constituents, showed better anticancer activity towards the cell of HepG2 cancer. [40] The X-linked inhibitor of apoptosis as a new molecular target for anticancer drugs used to resist the cancer cells to chemo and radiation therapy. [41] Phytochemical studies on active anti-colorectal cancer compounds *Alkanna tinctoria* and isolated eight quinone compounds. Among that alkannin, angelylalkannin, 5-methoxyangenyalkannin compounds show strong antiproliferative effects.

#### HIV

[42] This study shows enhance therapeutic options against AIDS by examining compounds of 4-thiazolidinone and its derivatives because of its antiviral action. [43] Another study, used to find lead molecules from the library as HIV-1 reverse transcriptase inhibitor.

#### Epilepsy

[44] This study deals with derived compounds of some herbal drugs and its quantitative structure-activity relationships done to found the inhibitory activity in epilepsy.

#### In skin

[45] This study found penetration activity of menthol which was found increase in temperature and concentration with a range between 1–20% increases menthols penetration increasing capacity.

[46] A study on on ginger and its 12 main components and human cytochrome P450.

[47] Use of medical herbs and its pharmacokinetic and dynamic effects against cytochrome P450.

[48] Interaction of Fangjifuling decoction, with CYP2D6 by virtual screening.

#### Contraceptive

[49] Piperine as a contraceptive in males.

## DISCUSSION

Bioinformatics is a research methodology in which using standard analytical techniques to find out action of complex mixture. Applying this, it is possible to decode the complex Ayurvedic formulations by using analytical tools like XRD, XRF, ICP-MS and the data from these can be processed to identify the structural formula. This structural formula can be used to assess the binding of the molecule to various proteins/receptors using standard database/software. From this review, we can identify that In-silico studies were more used in Chinese medicine. This study helps to identify the exact target site and all the above studies undergone the in-vitro, in-vivo studies later. Many studies are found in Traditional Chinese medicine to propagate the herbal medicine in china. [50] A Study found to understand the molecular mechanisms of TCM. [51] Another study reveals that usage of Traditional Chinese Medicine instead of conventional medicine. To study and explore TCM Quantitative information is important. TCM-ID gives information about number of prescription as 1588, herbs- 1313 herbal ingredients-5669, and 3725 herbal ingredients by its 3D structure. [52] Herb-drug interaction happened by enzymatic induction and inhibition. This interaction can be affect the circulation of drug and changes the clinical result.

## CONCLUSION

Bioinformatics is an approach which can be used to design novel combinations by first validating the concept on a computer and validating scenario. It solves many problems related to drug standardization, safety, and efficacy of the medicines. The latest technology which helps to reduce the time and resources required for an in- depth research in herbal formulations and *Rasaushadhis*. It is possible to assess particular drug with specific disease cell without any clinical trial. Bioinformatics-as a methodology in preclinical analysis of *Rasaushadhi* helps to perform the in-vivo studies in future studies.

**Source of Support:** Nil.

**Conflict of Interest:** None Declared.

## REFERENCES

1. Hong C, Cao J, Wu C-F, *et al.* The Chinese herbal formula Free and Easy Wanderer ameliorates oxidative stress through KEAP1-NRF2/HO-1 pathway. *Sci Rep.* 2017; 7(1):11551. doi:10.1038/s41598-017-10443-6
2. Hong M, Li S, Wang N, Tan H-Y, Cheung F, Feng Y. A Biomedical Investigation of the Hepatoprotective Effect of *Radix salviae miltiorrhizae* and Network Pharmacology-Based Prediction of the Active Compounds and Molecular Targets. *Int J Mol Sci* 2017; 18(3):620. doi:10.3390/ijms18030620
3. Yi F, Sun L, Xu L-J, *et al.* In silico Approach for Anti-Thrombosis Drug Discovery: P2Y1R Structure-Based TCMs Screening. *Front Pharmacol.* 2016; 7:531. doi:10.3389/fphar.2016.00531
4. Maione F, Cantone V, Chini MG, De Feo V, Mascolo N, Bifulco G. Molecular mechanism of tanshinone IIA and cryptotanshinone in platelet anti-aggregating effects: an integrated study of pharmacology and computational analysis. *Fitoterapia.* 2015; 100:174-178. doi:10.1016/j.fitote.2014.11.024
5. Han X-D, Zhou Z-W, Yang W, *et al.* A computational and functional study elicits the ameliorating effect of the Chinese herbal formula *Huo Luo Xiao Ling Dan* on experimental ischemia-induced myocardial injury in rats via inhibition of apoptosis. *Drug Des Devel Ther.* 2015; 9:1063-1102. doi:10.2147/DDDT.S76336
6. Jiang B, Chen J, Xu L, *et al.* Salvianolic acid B functioned as a competitive inhibitor of matrix metalloproteinase-9 and efficiently prevented cardiac remodeling. *BMC Pharmacol.* 2010; 10(1):10. doi:10.1186/1471-2210-10-10
7. Luo G, Lu F, Qiao L, Chen X, Li G, Zhang Y. Discovery of Potential Inhibitors of Aldosterone Synthase from Chinese Herbs Using Pharmacophore Modeling, Molecular Docking, and Molecular Dynamics Simulation Studies. *Biomed Res Int.* 2016; 2016:4182595. doi:10.1155/2016/4182595
8. Rahayu B, Baktiyani SCW, Nurdiana N. Theobroma cacao increases cells viability and reduces IL-6 and sVCAM-1 level in endothelial cells induced by plasma from preeclamptic patients. *Pregnancy Hypertens.* 2016; 6(1):42-46. doi:10.1016/j.preghy.2016.01.001
9. Kuok C-F, Hoi S-O, Hoi C-F, *et al.* Synergistic antibacterial effects of herbal extracts and antibiotics on methicillin-resistant *Staphylococcus aureus*: A computational and experimental study. *Exp Biol Med (Maywood).* 2017; 242(7):731-743. doi:10.1177/1535370216689828
10. Ding W, Gu J, Cao L, *et al.* Traditional Chinese herbs as chemical resource library for drug discovery of anti-infective and anti-inflammatory. *J Ethnopharmacol* 2014; 155(1):589-598. doi:10.1016/j.jep.2014.05.066
11. Kadioglu O, Jacob S, Bohnert S, *et al.* Evaluating ancient Egyptian prescriptions today: Anti-inflammatory activity of *Ziziphus spina-christi*. *Phytomedicine.* 2016; 23(3):293-306. doi:10.1016/j.phymed.2016.01.004
12. Shukla KL, Gund TM, Meshnick SR. Molecular modeling studies of the artemisinin (qinghaosu)-hemin interaction: docking between the antimalarial agent and its putative receptor. *J Mol Graph* 1995; 13(4):215-222. <http://www.ncbi.nlm.nih.gov/pubmed/8527414>. Accessed January 19, 2018.
13. Koushik Kumar G, Prasanna G, Marimuthu T, Saraswathi NT. Structural basis for complementary and alternative medicine: Phytochemical interaction with non-structural protein 2 protease-a reverse engineering strategy. *Chin J Integr Med* 2015; 21(6):445-452. doi:10.1007/s11655-014-1764-z
14. Skariyachan S, Prakash N, Bharadwaj N. In silico exploration of novel phytoligands against probable drug target of *Clostridium tetani*. *Interdiscip Sci.* 2012; 4(4):273-281. doi:10.1007/s12539-012-0138-5

15. Cui L, Wang Y, Liu Z, *et al.* Discovering New Acetylcholinesterase Inhibitors by Mining the Buzhongyiqi Decoction Recipe Data. *J Chem Inf Model.* 2015; 55(11):2455-2463. doi:10.1021/acs.jcim.5b00449
16. Qiao L-S, Zhang X-B, Jiang L, Zhang Y-L, Li G-Y. Identification of potential ACAT-2 selective inhibitors using pharmacophore, SVM and SVR from Chinese herbs. *Mol Divers.* 2016; 20(4):933-944. doi:10.1007/s11030-016-9684-9
17. Fong P, Tong HHY, Ng KH, Lao CK, Chong Cl, Chao CM. In silico prediction of prostaglandin D2 synthase inhibitors from herbal constituents for the treatment of hair loss. *J Ethnopharmacol* 2015; 175:470-480. doi:10.1016/j.jep.2015.10.005
18. Zheng Y, Wu J, Feng X, *et al.* In silico Analysis and Experimental Validation of Lignan Extracts from *Kadsura longipedunculata* for Potential 5-HT<sub>1A</sub>R Agonists. Taylor P, ed. *PLoS One.* 2015; 10(6):e0130055. doi:10.1371/journal.pone.0130055
19. Wu K-Z, Jiang Y-G, Zuo Y, Li A-X. Establishment of the method for screening the potential targets and effective components of huatuo reconstruction pill. *Interdiscip Sci.* 2014; 6(2):125-132. doi:10.1007/s12539-013-0056-1
20. Haiyu X, Yang S, Yanqiong Z, *et al.* Identification of key active constituents of Buchang Naoxintong capsules with therapeutic effects against ischemic stroke by using an integrative pharmacology-based approach. *Mol Biosyst.* 2016; 12(1):233-245. doi:10.1039/C5MB00460H
21. Mohd Fauzi F, John CM, Karunanidhi A, *et al.* Understanding the mode-of-action of *Cassia auriculata* via in silico and in vivo studies towards validating it as a long term therapy for type II diabetes. *J Ethnopharmacol* 2017; 197:61-72. doi:10.1016/j.jep.2016.07.058
22. Natarajan A, Sugumar S, Bitragunta S, Balasubramanyan N. Molecular docking studies of (4Z, 12Z)-cyclopentadeca-4, 12-dienone from *Grewia hirsuta* with some targets related to type 2 diabetes. *BMC Complement Altern Med.* 2015; 15(1):73. doi:10.1186/s12906-015-0588-5
23. Fayaz SM, Suvanish Kumar VS, Rajanikant KG. Finding needles in a haystack: application of network analysis and target enrichment studies for the identification of potential anti-diabetic phytochemicals. Ahmad M, ed. *PLoS One.* 2014; 9(11):e112911. doi:10.1371/journal.pone.0112911
24. Vaidya HB, Ahmed AA, Goyal RK, Cheema SK. Glycogen phosphorylase-a is a common target for anti-diabetic effect of iridoid and secoiridoid glycosides. *J Pharm Pharm Sci* 2013; 16(4):530-540. <http://www.ncbi.nlm.nih.gov/pubmed/24210061>. Accessed January 20, 2018.
25. Wang X-L, Jiao F-R, Yu M, *et al.* Constituents with potent  $\alpha$ -glucosidase inhibitory activity from *Pueraria lobata* (Willd.) ohwi. *Bioorg Med Chem Lett.* 2017; 27(9):1993-1998. doi:10.1016/j.bmcl.2017.03.013
26. Suhitha S, Devi SK, Gunasekaran K, Pakyntein HC, Bhattacharjee A, Velmurugan D. Phytochemical analyses and activity of herbal medicinal plants of North- East India for anti-diabetic, anti-cancer and anti-tuberculosis and their docking studies. *Curr Top Med Chem.* 2015; 15(1):21-36. <http://www.ncbi.nlm.nih.gov/pubmed/2579573>. Accessed January 20, 2018.
27. Alugolu V, Rentala S, Komarraju AL, Parimi UD. Docking studies of piperine - iron conjugate with human CYP450 3A4. *Bioinformation.* 2013; 9(7):334-338. doi:10.6026/97320630009334
28. Tomy MJ, Sharanya CS, Dileep K V, *et al.* Derivatives form better lipoxigenase inhibitors than piperine: in vitro and in silico study. *Chem Biol Drug Des.* 2015; 85(6):715-721. doi:10.1111/cbdd.12455
29. Wang S-Q, Du Q-S, Zhao K, Li A-X, Wei D-Q, Chou K-C. Virtual screening for finding natural inhibitor against cathepsin-L for SARS therapy. *Amino Acids.* 2007; 33(1):129-135. doi:10.1007/s00726-006-0403-1
30. Senol FS, Ślusarczyk S, Matkowski A, *et al.* Selective in vitro and in silico butyrylcholinesterase inhibitory activity of diterpenes and rosmarinic acid isolated from *Perovskia atriplicifolia* Benth. and *Salvia glutinosa* L. *Phytochemistry.* 2017; 133:33-44. doi:10.1016/j.phytochem.2016.10.012
31. Cui Z, Sheng Z, Yan X, Cao Z, Tang K. In Silico Insight into Potential Anti-Alzheimer's Disease Mechanisms of Icaritin. *Int J Mol Sci* 2016; 17(1):113. doi:10.3390/ijms17010113
32. Rajmohamed MA, Natarajan S, Palanisamy P, Abdulkader AM, Govindaraju A. Antioxidant and Cholinesterase Inhibitory Activities of Ethyl Acetate Extract of *Terminalia chebula*: Cell-free In vitro and In silico Studies. *Pharmacogn Mag.* 2017; 13(Suppl 3):S437-S445. doi:10.4103/pm.pm\_57\_17
33. Seo E-J, Saeed M, Law BYK, *et al.* Pharmacogenomics of Scopoletin in Tumor Cells. *Molecules.* 2016; 21(4):496. doi:10.3390/molecules21040496
34. Jayadeepa RM, Sharma S. Computational models for 5 $\alpha$ R inhibitors for treatment of prostate cancer: review of previous works and screening of natural inhibitors of 5 $\alpha$ R2. *Curr Comput Aided Drug Des.* 2011; 7(4):231-237. <http://www.ncbi.nlm.nih.gov/pubmed/22050680>. Accessed January 19, 2018.
35. Zhou X, Wang Y, Hu T, *et al.* Enzyme kinetic and molecular docking studies for the inhibitions of miltirone on major human cytochrome P450 isozymes. *Phytomedicine.* 2013; 20(3-4):367-374. doi:10.1016/j.phymed.2012.09.021
36. Jo AR, Kim JH, Yan X-T, Yang SY, Kim YH. Soluble epoxide hydrolase inhibitory components from *Rheum undulatum* and in silico approach. *J Enzyme Inhib Med Chem.* 2016; 31(sup2):70-78. doi:10.1080/14756366.2016.1189421
37. Shi X, Zhang G, Mackie B, Yang S, Wang J, Shan L. Comparison of the in vitro metabolism of psoralidin among different species and characterization of its inhibitory effect against UDP-glucuronosyltransferase (UGT) or cytochrome p450 (CYP450) enzymes. *J Chromatogr B* 2016; 1029-1030:145-156. doi:10.1016/j.jchromb.2016.06.031
38. Zhao G-F, Huang Z-A, Du X-K, Yang M-L, Huang D-D, Zhang S. Molecular docking studies of Traditional Chinese Medicinal compounds against known protein targets to treat non-small cell lung carcinomas. *Mol Med Rep.* 2016; 14(2):1132-1138. doi:10.3892/mmr.2016.5350
39. Chen L, Du J, Dai Q, Zhang H, Pang W, Hu J. Prediction of anti-tumor chemical probes of a traditional Chinese medicine formula by HPLC fingerprinting combined with molecular docking. *Eur J Med Chem.* 2014; 83:294-306. doi:10.1016/j.ejmech.2014.06.037
40. Nikolovska-Coleska Z, Xu L, Hu Z, *et al.* Discovery of embelin as a cell-permeable, small-molecular weight inhibitor of XIAP through structure-based computational screening of a traditional herbal medicine three-dimensional structure database. *J Med Chem* 2004; 47(10):2430-2440. doi:10.1021/jm030420+
41. Tung NH, Du G-J, Yuan C-S, Shoyama Y, Wang C-Z. Isolation and chemopreventive evaluation of novel naphthoquinone compounds from *Alkanna tinctoria*. *Anticancer Drugs.* 2013; 24(10):1058-1068. doi:10.1097/CAD.000000000000017
42. Seniya C, Yadav A, Khan GJ, Sah NK. In-silico Studies Show Potent Inhibition of HIV-1 Reverse Transcriptase Activity by a Herbal Drug. *IEEE/ACM Trans Comput Biol Bioinforma.* 2015; 12(6):1355-1364. doi:10.1109/TCBB.2015.2415771
43. Syahdi RR, Mun'im A, Suhartanto H, Yanuar A. Virtual screening of Indonesian herbal database as HIV-1 reverse transcriptase inhibitor. *Bioinformation.* 2012; 8(24):1206-1210. doi:10.6026/97320630081206
44. Mathew S, Faheem M, Al-Malki AL, Kumosani TA, Qadri I. In silico inhibition of GABARAP activity using antiepileptic medicinal derived compounds. *Bioinformation.* 2015; 11(4):189-195. doi:10.6026/97320630011189
45. Wan G, Dai X, Yin Q, Shi X, Qiao Y. Interaction of menthol with mixed-lipid bilayer of stratum corneum: A coarse-grained simulation study. *J Mol Graph Model* 2015; 60:98-107. doi:10.1016/j.jmgm.2015.06.005
46. Zhou S-F, Qiu J, Zhou Z-W, He Z, Zhang X, Zhu S. Estimation of the binding modes with important human cytochrome P450 enzymes, drug interaction potential, pharmacokinetics, and hepatotoxicity of ginger components using molecular docking, computational, and pharmacokinetic modeling studies. *Drug Des Devel Ther.* 2015; 9:841. doi:10.2147/DDDT.S74669
47. Zhou S, Gao Y, Jiang W, Huang M, Xu A, Paxton JW. Interactions of herbs with cytochrome P450. *Drug Metab Rev.* 2003; 35(1):35-98. doi:10.1081/DMR-120018248
48. Mo S-L, Liu W-F, Li C-G, *et al.* Pharmacophore, QSAR, and binding mode studies of substrates of human cytochrome P450 2D6 (CYP2D6) using molecular docking and virtual mutations and an application to chinese herbal medicine screening. *Curr Pharm Biotechnol.* 2012; 13(9):1640-1704. <http://www.ncbi.nlm.nih.gov/pubmed/22039821>. Accessed January 19, 2018.
49. Chinta G, Ramya Chandar Charles M, Klopčič I, Sollner Dolenc M, Periyasamy L, Selvaraj Coumar M. In Silico and In Vitro Investigation of the Piperine's Male Contraceptive Effect: Docking and Molecular Dynamics Simulation Studies in Androgen-Binding Protein and Androgen Receptor. *Planta Med.* 2015; 81(10):804-812. doi:10.1055/s-0035-1546082
50. Yan J, Wang Y, Luo S-J, Qiao Y-J. TCM grammar systems: an approach to aid the interpretation of the molecular interactions in Chinese herbal medicine. *J Ethnopharmacol.* 2011; 137(1):77-84. doi:10.1016/j.jep.2011.04.057
51. Ganguli S, Mazumder M, Basu P, Roy P, Mitra S, Datta A. In silico screening of herbal and nanoparticle lead compounds for effectivity against H5N1, H1N1 neuraminidase and telomerase. *BMC Bioinformatics.* 2010; 11 Suppl 1(Suppl 1):S13. doi:10.1186/1471-2105-11-S1-S13
52. Chen X-W, Sneed KB, Pan S-Y, *et al.* Herb-drug interactions and mechanistic and clinical considerations. *Curr Drug Metab.* 2012; 13(5):640-651. <http://www.ncbi.nlm.nih.gov/pubmed/22292789>. Accessed January 23, 2018.

**HOW TO CITE THIS ARTICLE**

Monisha M, Mishra AK, Ramesh NV, Pillai KU, Vineeth PK. *In-silico* studies in herbal drugs: A review. J Ayu Herb Med 2018;4(1):43-47.